

A generator system of invariant differential forms.

Tensai Bakabon

Bakada University, Department of Mathematics
3-10-5 Tsukimidai Susukigahara, Nerima-ku, Tokyo, Japan
e-mail:korede.ii.noda@yahoo.co.jp

April 1, 2002

Abstract

We obtain a generator system of the algebra of $\mathrm{GL}(V)$ -invariant differential forms on $\mathrm{End}_{\mathbf{k}}(V)$. The proof uses the Weyl-Schur reciprocity.

Let $\mathbf{k}$ be a field of characteristic zero and $M_n(\mathbf{k})$ be the set of all $n \times n$ matrices with entries of elements of $\mathbf{k}$. We consider $\mathrm{GL}_n(\mathbf{k})$ adjoint action on $M_n(\mathbf{k})$, i.e. for given $g \in \mathrm{GL}_n(\mathbf{k})$ and $X \in M_n(\mathbf{k})$, $\mathrm{ad}(g)X := g \cdot X \cdot g^{-1}$. It is classically known that the algebra $\mathbb{C}[M_n(\mathbf{k})]^{\mathrm{GL}_n(\mathbf{k})}$ of all $\mathrm{GL}_n(\mathbf{k})$ -invariant polynomials on $M_n(\mathbf{k})$ is generated by $\mathrm{Tr}(X^\ell)$ ($\ell = 1, 2, \dots$), where $X = (x_{ij})_{i,j=1}^n$. More precisely, if we define polynomials $\sigma_1(X), \dots, \sigma_n(X)$ of X by

$$\det(t \cdot I_n + X) = t^n + \sigma_1(X)t^{n-1} + \dots + \sigma_n(X),$$

the invariant ring is the polynomial ring generated by $\sigma_1(X), \dots, \sigma_n(X)$ over $\mathbf{k}$. Furthermore polynomials $\sigma_1(X), \dots, \sigma_n(X)$ and $\mathrm{Tr}(X^\ell)$ ($\ell = 1, 2, \dots$) are connected by Newton's formula.

We next consider $\mathrm{GL}_n(\mathbf{k})$ -invariant differential forms on $M_n(\mathbf{k})$ with polynomial coefficients. Similar to the above case, we obtain many invariant differential forms of degree p by the formula

$$\begin{aligned} \omega^{\ell_1 \ell_2 \dots \ell_p} &:= \mathrm{Tr} \left(X^{\ell_1} dX \wedge X^{\ell_2} dX \wedge \dots \wedge X^{\ell_p} dX \right) \\ &= \sum_{i_1, \dots, i_p, j_1, \dots, j_p=1}^n (X^{\ell_1})_{i_1 j_1} dx_{j_1 i_2} \wedge (X^{\ell_2})_{i_2 j_2} dx_{j_2 i_3} \dots (X^{\ell_p})_{i_p j_p} dx_{j_p i_1}, \end{aligned}$$

where $(X^\ell)_{ij}$ is the (i, j) component of X^ℓ and $dX = (dx_{ij})_{i,j=1}^n$.

It is natural to ask if the algebra of all $\mathrm{GL}_n(\mathbf{k})$ -invariant differential forms is generated by these forms. The purpose of this paper is to answer this question affirmatively.

Theorem 1 *The algebra of all $\mathrm{GL}_n(\mathbf{k})$ -invariant differential forms on $M_n(\mathbf{k})$ is generated by $\omega^{\ell_1 \ell_2 \dots \ell_p} = \mathrm{Tr}(X^{\ell_1} dX \wedge X^{\ell_2} \dots \wedge X^{\ell_p} dX)$, $(p \geq 0, \ell_1, \dots, \ell_p \geq 0)$ over $\mathbf{k}$. In other words, any invariant form is expressed as a linear combination of forms of the above type and their product.*

Proof of Theorem.

Put $V = \mathbf{k}^n$, $E := \mathrm{End}_{\mathbf{k}}(V) \cong M_n(\mathbf{k}) \cong V \otimes V^*$ and $G := \mathrm{GL}(V)$. Then we can identify E and its dual E^* because $E \ni X, Y \mapsto \mathrm{Tr}(XY)$ is a $\mathrm{GL}(V)$ -invariant non-degenerate bilinear pairing. So the set of all differential forms on E is

$$\Omega(E) \cong \mathcal{S}(E) \otimes \Lambda(E) \cong \bigoplus_{p,q \geq 0} \mathcal{S}^p(E) \otimes \Lambda^q(E)$$

where $\mathcal{S}(E)$ and $\Lambda(E)$, respectively, are symmetric and alternative tensor algebra of the vector space E . Hence it is sufficient to show that each component $\mathcal{S}^p(E) \otimes \Lambda^q(E)$ is generated by $\omega^{\ell_1 \ell_2 \dots \ell_r}$ ($\ell_1 + \dots + \ell_r \leq p, r \leq q$). Consider the G -equivariant projection map $\mathcal{P} : E^{\otimes(p+q)} \rightarrow \mathcal{S}^p(E) \otimes \Lambda^q(E)$

$$\begin{array}{ccc} E^{\otimes(p+q)} & \ni & v_1 \otimes \dots \otimes v_p \otimes w_1 \otimes \dots \otimes w_q \\ \downarrow & & \downarrow \\ \mathcal{S}^p(E) \otimes \Lambda^q(E) & \ni & \frac{1}{p!q!} \sum_{\sigma \in \mathfrak{S}_p, \tau \in \mathfrak{S}_q} \mathrm{sgn}(\tau) v_{\sigma^{-1}(1)} \otimes \dots \otimes v_{\sigma^{-1}(p)} \otimes w_{\tau^{-1}(1)} \otimes \dots \otimes w_{\tau^{-1}(q)}. \end{array}$$

Since G is a reductive group, $\mathcal{P}$ induces a surjection of G -invariant tensors to G -invariant differential forms.

$$\mathcal{P}|_{(E^{\otimes(p+q)})^G} : (E^{\otimes(p+q)})^G \twoheadrightarrow (\mathcal{S}^p(E) \otimes \Lambda^q(E))^G.$$

Thus we reduce the problem to description of G -invariant tensors and its images by $\mathcal{P}$. Recall $E^{\otimes r} \cong \mathrm{End}_{\mathbf{k}}(V^{\otimes r})$ and $(E^{\otimes r})^G \cong \mathrm{End}_G(V^{\otimes r})$, we have a special element $\Phi(\rho)$ of G -invariant tensors $(E^{\otimes r})^G$ corresponding to a permutation $\rho \in \mathfrak{S}_r$,

$$\mathrm{End}_G(V^{\otimes r}) \ni \Phi(\rho) : e_1 \otimes \dots \otimes e_r \mapsto e_{\rho^{-1}(1)} \otimes \dots \otimes e_{\rho^{-1}(r)}.$$

Recall that the Weyl-Schur reciprocity assert that

$$\{\Phi(\rho) \in (E^{\otimes r})^G \mid \rho \in \mathfrak{S}_r\} \text{ form a basis of } (E^{\otimes r})^G.$$

Put $r = p + q$, it is easily checked that $\mathcal{P}(\Phi(\rho)) \in (\mathcal{S}^p(E) \otimes \Lambda^q(E))^G$ is a product of $\omega^{\ell_1 \ell_2 \cdots \ell_s}$ for appropriate s and $\ell_1, \ell_2, \dots, \ell_s$. In particular, the number of factors of $\mathcal{P}(\Phi(\rho))$ is equal to the number of cyclic permutations in the cyclic decomposition of $\rho \in \mathfrak{S}_r$. This completes the proof.

□

Perhaps it helps some calculations in relative de Rham cohomologies of invariant maps [Ve].

Acknowledgement

The author thanks Ms.Ys.(real) for decisive help in the proof of our main result.

References

- [GW] R. Goodman, N. R. Wallach, Representations and invariants of the classical groups. Cambridge.
- [Ve] J. Vey, Un problème de cohomologie relative. Ark. Mat. 15 (1977), no. 1, 21–31.